INORGANICA CHIMICA ACTA VOLUME 198–200, AUGUST–OCTOBER, 1992

Contents

Preface xi

1991 Publication Statistics xii

Articles

Have adenosine 5′-triphosphate (ATP^{4-}) and related purine-nucleotides played a role in early evolution? ATP, its own 'enzyme' in metal ion facilitated hydrolysis! 1
H. Sigel (Basel, Switzerland)

Crystallographic disorder in $[M_2X_8]^{n-}$, $M_2X_4L_4$ and related compounds: chemical and theoretical implications 13
F. A. Cotton and J. L. Eglin (College Station, TX, USA)

The preparation of new manganese(II) isocyanide complexes, $MnI_2(CNBu^t)_n$ ($n=1$, 1.5, 2) and the mixed isocyanide/tertiary-phosphine complex $MnI_2(PPh_3)(CNBu^t)_2$. The isolation and X-ray crystallographic characterisation of the Mn^I/Mn^{II} mixed-valence isomeric complexes $[Mn(CNBu^t)_6][MnI_3(PPh_3)]$ and $[Mn(CNBu^t)_5(PPh_3)][MnI_3(CNBu^t)]$ 23
S. M. Godfrey, G. Q. Li, C. A. McAuliffe, P. T. Ndifon and R. G. Pritchard (Manchester, UK)

Dioxygen uptake and transfer by Co(III), Rh(III) and Ir(III) catecholate complexes 31
P. Barbaro, C. Bianchini, K. Linn, C. Mealli, A. Meli, F. Vizza (Florence, Italy), F. Laschi and P. Zanello (Siena, Italy)

α- versus β-Hydrogen elimination in the formation of propene from an osmacyclobutane 57
W. Fischer, R. T. Hembre, D. R. Sidler and J. R. Norton (Fort Collins, CO, USA)

Oxygen atom transfer in the reaction between hexakis(dimethyl-tert-butylsiloxy)ditungsten and nitric oxide. A remarkable difference in the reactivity of the tungsten–tungsten triple bond as a function of the attendant ligands: Bu^tO versus Bu^tMe_2SiO 63
M. H. Chisholm, C. M. Cook, K. Folting and W. E. Streib (Bloomington, IN, USA)

Synthesis and antitumor activity of water-soluble 2-benzyl-1,2-diaminobutane-α-oxycarboxylatoplatinum(II) complexes 79
H. Brunner, F. Maiterth and B. Treittinger (Regensburg, FRG)

Ni^{IV} cubanes: synthesis and characterization of the $[Ni_4Se_4(Se_3)_5(Se_4)]^{4-}$ and $[Ni_4Te_4(Te_2)_2(Te_3)_4]^{4-}$ anions 85
J. M. McConnachie, M. A. Ansari and J. A. Ibers (Evanston, IL, USA)

Structural investigations of the hexavanadium core $\{V_6O_{19}\}$ in 'oxidized', mixed valence and 'reduced' clusters of the type $[V^V_{6-n}V^{IV}_nO_{13-n}(OH)_n\{(OCH_2)_3CR\}_2]^{2-}$, $n=0$, 3 and 6 95
Q. Chen and J. Zubieta (Syracuse, NY, USA)

An investigation of the reaction of $[Rh_2(COMe)_2(CO)_2I_6]^{2-}$ with some tertiary phosphines, and $PPh(OMe)_2$, and the X-ray structure of $[Rh(COMe)I_2(Ph_2PCH_2PPh_2)]$ 111
H. Adams, N. A. Bailey, B. E. Mann and C. P. Manuel (Sheffield, UK)

One- and two-dimensional rare earth–copper molecular materials 119
O. Guillou, O. Kahn, R. L. Oushoorn, K. Boubekeur and P. Batail (Orsay, France)

On the structures, stabilities and fragmentation patterns of carbon clusters including Buckminsterfullerene 133
J. J. Novoa, M.-H. Whangbo (Raleigh, NC, USA) and J. M. Williams (Argonne, IL, USA)

Synthesis and crystal structure of a molybdenum carbonyl complex, $Mo(CO)_4[cyclo(Ph_6P_3As_3)]$, with scrambled phosphorus/arsenic identities obtained from a perfectly alternating phosphorus–arsenic heterocycle, 1,3,5-tris(phenylphospha)-2,4,6-tris(phenylarsa)cyclohexane, $[cyclo(Ph_6P_3As_3)]$ 139
A. L. Rheingold and F. P. Arnold (Newark, DE, USA)

Electronic absorption spectra of M(II)(Met121X) azurins (M=Co, Ni, Cu; X=Leu, Gly, Asp, Glu): charge-transfer energies and reduction potentials 145
A. J. Di Bilio, T. K. Chang, B. G. Malmström, H. B. Gray (Pasadena, CA, USA), B. G. Karlsson, M. Nordling, T. Pascher and L. G. Lundberg (Göteborg, Sweden)

The synthesis of *endo*-3-diphenylphosphino-(1*R*)-(+)-camphor (L) and some of its complexes with palladium(II), platinum(II) and rhodium(I); crystal structures of L and *cis*-$[PdCl_2L_2]$ 149
S. D. Perera, B. L. Shaw, M. Thornton-Pett and J. D. Vessey (Leeds, UK)

Elsevier Sequoia

Monocomplex formation and dissociation of some first row divalent transition metal ions with 2-chloro-1,10-phenanthroline by the high-pressure stopped-flow technique 159
G. Laurenczy, P. Bugnon and A. E. Merbach (Lausanne, Switzerland)

Synthesis and structures of novel types of heteronuclear Pt–M (M = Ag, Sn or Pb) neutral or anionic organometallic complexes 165
R. Usón and J. Forniés (Saragossa, Spain)

Coordination chemistry of nitriles and cyanamide at electron-rich metal centres 179
A. J. L. Pombeiro (Lisbon, Portugal)

Mechanistic information on the reaction of model palladium(II) complexes with purine nucleosides and 5′-nucleotides in reference to the antitumor activity of related platinum complexes 187
M. Shoukry, H. Hohmann and R. van Eldik (Witten, FRG)

Activation of dithioformate by iron complexes: insertion of alkynes into the coordinated C–S bond. X-ray structures of $\overline{Fe[\eta^3\text{-}H\text{–}B(H)_2 \leftarrow S\text{–}C(SMe)H]}(CO)(PMe_3)_2$ and $\overline{Fe[\eta^3\text{-}S\text{–}C(Ph)=CH\text{–}CH(SMe)]}(CO)(PMe_3)_2$ 193
D. V. Khasnis, N. Pirio, D. Touchard, L. Toupet and P. H. Dixneuf (Rennes, France)

Subvalent Group 14 metal compounds
XIV. The X-ray crystal structures of two monomeric Group 14 metal bisamides, $Ge[N(SiMe_3)_2]_2$ and $Sn[\overline{NC(Me)_2(CH_2)_3CMe_2}]_2$ 203
R. W. Chorley, P. B. Hitchcock, M. F. Lappert, W.-P. Leung (Brighton, UK), P. P. Power and M. M. Olmstead (Davis, CA, USA)

The 1,3-dimetalloallyl tri-ruthenium cluster $(\mu\text{-}H)Ru_3(\mu_3\text{-}\eta^3\text{-}CHCHCMe)(CO)_9$ and its PPh_3 derivatives 211
K. M. Rao, R. J. Angelici and V. G. Young, Jr. (Ames, IA, USA)

Kinetics and mechanism of CO ligand substitution in the ring-substituted indenyl rhodium complexes $[(\eta^5\text{-}C_9R_nH_{7-n})Rh(CO)_2]$ 219
A. K. Kakkar, N. J. Taylor, T. B. Marder (Waterloo, Ont., Canada), J. K. Shen, N. Hallinan and F. Basolo (Evanston, IL, USA)

Axial ligand bonding in blue copper proteins 233
M. D. Lowery and E. I. Solomon (Stanford, CA, USA)

Mass spectroscopic evidence for the formation of mixed-metal octahedral clusters $[Mo_nW_{6-n}Cl_{14}]^{2-}$ 245
H. A. Hodali, H. Hung and D. F. Shriver (Evanston, IL, USA)

Organotin(IV) compounds of 2-thiopyridine. Crystal and molecular structure of dicyclohexyltin(IV) bis(2-pyridyl-thiolate) 249
M. Bouâlam (Brussels, Belgium and Tétouan, Moroccco), J. Meunier-Piret (Louvain-la-Neuve, Belgium), M. Biesemans, R. Willem and M. Gielen (Brussels, Belgium)

Phosphorus–carbon bond cleavage at a di-iron centre: synthesis of μ-phosphidomethyl complexes $[Fe_2(CO)_6(\mu\text{-}CH_2PR_2)(\mu\text{-}PR_2)]$ from $[Fe_2(CO)_6(\mu\text{-}CO)(\mu\text{-}R_2PCH_2PR_2)]$ 257
N. M. Doherty, G. Hogarth, S. A. R. Knox, K. A. Macpherson, F. Melchior, D. A. V. Morton and A. G. Orpen (Bristol, UK)

Use of the 2,6-bis[(dimethylamino)methyl]phenyl ligand to form some pentacoordinate derivatives of P(III), As(III), Sb(III) and Bi(III) 271
D. A. Atwood, A. H. Cowley and J. Ruiz (Austin, TX, USA)

Synthesis and NMR structural studies of allyl(polypyrazolylborate)palladium and platinum complexes 275
K. Ohkita, H. Kurosawa, T. Hasegawa, T. Shirafuji and I. Ikeda (Osaka, Japan)

Synthesis, characterization and reactivity of some mono- and dinuclear chlororuthenium complexes containing chelating ditertiary phosphines (P–P) with P–P:Ru = 1 283
A. M. Joshi, I. S. Thorburn, S. J. Rettig and B. R. James (Vancouver, BC, Canada)

The reactivity of spectroscopically detected peroxy complexes of iron porphyrins 297
A. L. Balch (Davis, CA, USA)

Beryllocene and related slip-sandwich structures 309
J. K. Beattie and K. W. Nugent (Sydney, NSW, Australia)

New trends in polyoxometalate chemistry, toward large polyanions, toward nitrosyl-substituted polyanions 319
Y. Jeannin, G. Hervé and A. Proust (Paris, France)

Preparation and reactions of some organosilicon cyanates 337
C. Eaborn, Y. Y. El-Kaddar and P. D. Lickiss (Brighton, UK)

Application of the ligand polyhedral model to dicobalt octacarbonyl 345
B. F. G. Johnson and E. Parisini (Edinburgh, Scotland, UK)

Transition-metal complexes with sulfur ligands
Part LXXXIV. CH_3 complexes with [NiS] fragments as models for CO dehydrogenase. Synthesis and properties of $[Ni('MeS_2')_2]$ and $(NMe_4)[Ni(CH_3)('MeS_2')_2]$ ('MeS_2'$^-$ = *o*-(methylthio)thiophenolate(1−)) 351
D. Sellmann, H. Schillinger, F. Knoch and M. Moll (Erlangen, FRG)

A classification scheme for homogeneous metal catalyzed oxidations by O_2 359
R. S. Drago and R. H. Beer (Gainesville, FL, USA)

Anionic dicarbollyl complexes of germanium(II) and tin(II) — missing links in a series of π-complexes with Group 14 elements as central atoms 369
P. Jutzi, D. Wegener, H.-G. Stammler (Bielefeld, FRG), A. Karaulov and M. B. Hursthouse (Cardiff, UK)

Synthesis and comparative chemistry of the early–late transition metal heterobimetallacycles $CpCp'Ta(CH_2)_2Ir(CO)(L)$ and their main-group element–late transition metal analogues $R_2P(CH_2)_2Ir(CO)(L)$ 377
M. J. Hostetler, M. D. Butts and R. G. Bergman (Berkelely, CA, USA)

Synthesis and characterisation of some hexanuclear heterometallic clusters containing arene ligands; the crystal and molecular structures of $RuOs_5(CO)_{15}(\eta^6\text{-}C_6H_6)$ and $RhOs_5(CO)_{15}(\eta^5\text{-}C_5Me_5)$ 393
R. K. Henderson, P. A. Jackson, B. F. G. Johnson, J. Lewis and P. R. Raithby (Cambridge, UK)

Iron(II) thiolates as reversible carbon monoxide carriers 401
A. Szakács-Schmidt, J. Kreisz, L. Markó, Z. Nagy-Magos and J. Takács (Vezsprém, Hungary)

Electron impact (EI) and fast-atom bombardment (FAB) mass spectra of coordination complexes: aggregation processes in the mass spectra of metal derivatives of 1,2,3,4,5-pentakis(methoxycarbonyl)cyclopentadiene 407
M. I. Bruce and M. J. Liddell (Adelaide, SA, Australia)

Metal complexes of biologically important ligands
Part LXIV. Pentamethylcyclopentadienyl–rhodium(III) complexes with α-amino acid derivatives as bridging ligands and with several chiral metal centers: synthesis and structure of $[Cp^*Rh(\mu\text{-L-phenylalaninato})]_3(BF_4)_3$ and $[Cp^*Rh(\mu\text{-}NHCOCH_2NCO_2CH_2Ph)]_2$ 415
R. Krämer, K. Polborn, C. Robl and W. Beck (Munich, FRG)

New synthetic, selective, high-affinity ligands for effective trivalent metal ion binding and transport 421
R. J. Motekaitis, Y. Sun and A. E. Martell (College Station, TX, USA)

Synthesis and characterization of some new complexes of the Vaska-type *trans*-$[IrX(CO)L_2]$ (X = Cl or I; L = phosphite, phosphinite or phosphonite) and the direction of dihydrogen addition to these species 429
X.-L. Luo, D. Michos, R. H. Crabtree (New Haven, CT, USA) and M. B. Hall (College Station, TX, USA)

Ruthenium(II) complexes of a 2,2′-biphosphinine 437
D. Carmichael, P. Le Floch, L. Ricard and F. Mathey (Palaiseau, France)

Synthesis and characterization of compounds with boron–boron double bonds 443
P. P. Power (Davis, CA, USA)

Tris(1,2-ethanediamine) complexes of osmium(IV), osmium(III) and osmium(II): oxidative dehydrogenation reactions 449
P. A. Lay (Sydney, NSW, Australia) and A. M. Sargeson (Canberra, ACT, Australia)

Synthesis and reactivity of digermylplatinum(II) complexes $Me_2\overline{Ge(CH_2)_nGe(Me_2)}PtL_2$ (n = 0, 1, 2) 461
J. Barrau, G. Rima, V. Cassano and J. Satgé (Toulouse, France)

New trinuclear and hexanuclear ruthenium clusters derived from the reaction of $[Ru_3(CO)_{12}]$ with phenylphosphine or cyclohexylphosphine 469
A. J. Deeming, S. Doherty and N. I. Powell (London, UK)

NMR is a unique and necessary step in the investigation of iron sulfur proteins: the HiPIP from *R. gelatinosus* as an example 483
I. Bertini (Florence, Italy), F. Capozzi, C. Luchinat (Bologna, Italy), M. Piccioli and M. Vicens Oliver (Florence, Italy)

Molecular architecture of copper(I) thiometallate complexes. Example of a cubane with an extra face, $(NPr_4)_3[MS_4Cu_4Cl_5]$ (M = Mo, W) 493
Y. Jeannin, F. Sécheresse, S. Bernès and F. Robert (Paris, France)

Hexanuclear polypyridine complexes containing different metals, bridging ligands and/or terminal ligands. Absorption spectra, electrochemical oxidation, luminescence properties and intercomponent energy transfer. 507
G. Denti, S. Serroni (Pisa, Italy), S. Campagna, V. Ricevuto (Messina, Italy), A. Juris, M. Ciano and V. Balzani (Bologna, Italy)

Octahedro octahedra and tetrahedra
II. Tetra- to heptanuclear carbonyl(cyano)chromato, -molybdato and -tungstato complexes of 3d metals 513
M. Fritz, D. Rieger, E. Bär, G. Beck, J. Fuchs, G. Holzmann and W. P. Fehlhammer (Berlin, FRG)

Highly fluorinated tetrazanes and other polynitrogen compounds. 527
G. Sarwar, N. R. Patel, Y. Y. Zheng, E. O. John, R. L. Kirchmeier and J. M. Shreeve (Moscow, ID, USA)

Manganese(II)cyclopentadienide and cyclopentadienylmanganese(biphenyl) as starting materials for the synthesis of carbonyl free organomanganese complexes. 533
K. Jonas, C.-C. Häselhoff, R. Goddard and C. Krüger (Mülheim an der Ruhr, FRG)

Protonation of cobalt tetraneopentoxyphthalocyanine as a function of oxidation state 543
P. A. Bernstein and A. B. P. Lever (North York, Ont., Canada)

Metal insertion into a CH bond as a route to the heterobimetallic μ-methylidene complex $C_5H_5(CO)_2Re(\mu\text{-}CH_2)Pt(PPh_3)_2$ 557
C. P. Casey, Y. Wang, L. M. Petrovich, T. L. Underiner, P. N. Hazin and J. M. Desper (Madison, WI, USA)

Synthesis and reactivity of molybdenum–sulfur cubes. 565
C. L. Coyle, K. A. Eriksen, S. Farina, J. Francis, Y. Gea, M. A. Greaney, P. J. Guzi, T. R. Halbert, H. H. Murray and E. I. Stiefel (Annandale, NJ, USA)

Acyclic mononuclear and macrocyclic dinuclear silver(I) complexes of Schiff base ligands derived from *N*,*N*-bis(2-aminoalkyl)-2-phenylethylamines. 577
D. E. Fenton and P. C. Hellier (Sheffield, UK)

Preparation and spectroscopic characterization of (η^6-arene)ruthenium(II) trifluoromethylsulfonates and of dinuclear di-μ- and tri-μ-hydrido (η^6-arene)ruthenium(II) complexes. Homogeneous hydrogenation of cyclohexene and 1-hexene catalysed by various (η^6-arene)ruthenium(II) complexes 583
M. A. Bennett and J. P. Ennett (Canberra, ACT, Australia)

Synthesis of complexes with molybdenum–platinum bonds from the alkylidyne(carborane)molybdenum reagent $[NEt_4][Mo(\equiv CC_6H_4Me\text{-}4)(CO)_2(\eta^5\text{-}C_2B_9H_9Me_2)]$ 593
J. C. Jeffery, M. D. Mortimer and F. G. A. Stone (Bristol, UK)

Nucleophilic addition of chromium carbene anions to the coordinated olefin in $[Cp(CO)_2Fe(\eta^2\text{-olefin})]^+$ complexes and to the cyclohexadienyl ligand in $[(\eta^5\text{-cyclohexadienyl})(CO)_3Fe]^+$ 601
C. Kelley, M. R. Terry, A. W. Kaplan, G. L. Geoffroy (University Park, PA, USA), N. Lugan, R. Mathieu (Toulouse, France), B. S. Haggerty and A. L. Rheingold (Newark, DE, USA)

Magnetochemistry of the tetrahaloferrate(III) ions
6. Crystal structure and magnetic ordering in $[(pyH)_3Cl][FeCl_4]_2$ 613
R. Shaviv, C. B. Lowe (Chicago, IL, USA), J. A. Zora, C. B. Aakeröy, P. B. Hitchcock, K. R. Seddon (Brighton, UK) and R. L. Carlin (Chicago, IL, USA)

Procedures for the conversion of $\mu_2S \rightarrow \mu_2O$ in trinuclear $Mo(IV)_3$ incomplete cuboidal Mo/S cluster complexes. . . . 623
Q.-T. Liu, J. Lu and A. G. Sykes (Newcastle upon Tyne, UK)

Models of the oxidized forms of polyiron oxo proteins: synthetic routes to (μ-oxo)bis(μ-carboxylato)diiron(III) complexes with neutral monodentate and bidentate capping ligands 627
K. L. Taft, A. Masschelein, S. Liu, S. J. Lippard (Cambridge, MA, USA), D. Garfinkel-Shweky and A. Bino (Jerusalem, Israel)

Kinetics and mechanism of oxidative cleavage of the metal–metal bond in $[M_2(CO)_{10}]^{2-}$, M = Cr, Mo and W 633
J. R. Phillips and W. C. Trogler (La Jolla, CA, USA)

Heterometallic species containing Cp_2MH_2 (M = Mo and W) and Cu^+ or Ag^+ are inner sphere redox intermediates 639
L. F. Rhodes, J. C. Huffman and K. G. Caulton (Bloomington, IN, USA)

Spectroscopic and electrochemical aspects of copper(II) chelates containing 1,2-diamines and β-ketoenols 651
C. Tsiamis (Salonica, Greece)

Zinc complexes of cyclohexane triamine ligands 663
U. Brand and H. Vahrenkamp (Freiburg, FRG)

Characterisation of intermediates involved in the *mer*-$W(CO)_3(\eta^1\text{-dpm})(\eta^2\text{-dpm})$ to *mer*-$[W(CO)_2(\eta^2\text{-dpm})_2]^{2+}$ (dpm = $Ph_2PCH_2PPh_2$) oxidation process in acetone by steady-state and fast scan voltammetry at platinum and mercury microelectrodes 671
A. M. Bond, R. Colton (Bundoora, Australia) and R. S. Hutton (Parkville, Australia)

Structures and dynamic features of dinuclear tartratozirconocene complexes 679
G. Erker, M. Rump (Münster, FRG), C. Krüger and M. Nolte (Mülheim an der Ruhr, FRG)

Monoazadienes ruthenium carbonyl chemistry 689
C. J. Elsevier, W. P. Mul and K. Vrieze (Amsterdan, Netherlands)

Hydridosilicates: a new class of pentacoordinated silicon derivatives with unusual properties 705
R. Corriu, C. Guérin, B. Henner and Q. Wang (Montpellier, France)

Homo- and heterometallic transition-metal complexes containing bis(cyclopentadienyl)methane as ligand 715
H. Werner (Würzburg, FRG)

Metal-stabilized rare tautomers of nucleobases
4. On the question of adenine tautomerization by a coordinated platinum(II) 723
B. Lippert (Dortmund, FRG), H. Schöllhorn and U. Thewalt (Ulm, FRG)

The reaction of disilenes with P_4 and As_4 733
A. D. Fanta, R. P. Tan, N. M. Comerlato, M. Driess, D. R. Powell and R. West (Madison, WI, USA)

Intermediates of the cyclotrimerization of 2-butyne with a chromium catalyst 741
G. Wilke, H. Benn, R. Goddard, C. Krüger and B. Pfeil (Mülheim an der Ruhr, FRG)

Weak interactions in metal complexes of amino acids with a phosphorylated side chain. Conversion of aromatic ring stacking to electrostatic bonding by tyrosine phosphorylation 749
O. Yamauchi, A. Odani and H. Masuda (Nagoya, Japan)

New silanediyl (silylene) complexes of ruthenium(0) 763
H. Handwerker, C. Leis, S. Gamper and C. Zybill (Garching, FRG)

Synthesis and molecular structure of optically active oxazolidine complexes of platinum(II) and palladium(II) 771
A. Albinati, F. Lianza (Milan, Italy), H. Berger, C. Arz and P. S. Pregosin (Zurich, Switzerland)

Chemical hardness and the electronic chemical potential 781
R. G. Pearson (Santa Barbara, CA, USA)

The redox chemistry of the triply bonded compounds $Re_2X_4(\mu\text{-dmpm})_3$ (X = Cl or Br; dmpm = $Me_2PCH_2PMe_2$) and the isolation and characterization of the nitrosyl complex $Re_2Cl_5(\mu\text{-dmpm})_2(NO)$ 787
I. Ara and R. A. Walton (West Lafayette, IN, USA)

Strong exchange coupling between the lanthanide ions and the phthalocyaninato ligand radical in bis(phthalocyaninato)lanthanide sandwich compounds 795
K. L. Trojan, J. L. Kendall, K. D. Kepler and W. E. Hatfield (Chapel Hill, NC, USA)

^{199}Hg NMR investigation on the solution structure of Hg(II) complexes of oligopeptides containing cysteine and histidine residues 805
H. Adachi, N. Ueyama and A. Nakamura (Osaka, Japan)

Magnetic and spectral properties of paramagnetic metal-ion polyoxolene radical complexes 813
A. Dei and D. Gatteschi (Florence, Italy)

A Marcus-type model for steric and electronic effects of ligands in atom transfer reactions of metal carbonyl radicals 823
M.-G. Choi and T. L. Brown (Urbana-Champaign, IL, USA)

Synthesis and photophysical characterization of highly luminescent complexes of Ru(II) containing 4,4′-di-(*p*-carboxyphenyl)-2,2′-bipyridine 831
K. Kalyanasundaram, Md. K. Nazeeruddin, M. Grätzel (Lausanne, Switzerland), G. Viscardi, P. Savarino and E. Barni (Turin, Italy)

The icosahedron in inorganic chemistry. 841
R. B. King (Athens, GA, USA)

Covalent linkage of glucose oxidase to modified basal plane pyrolytic graphite electrodes and the use in the ferrocene-mediated amperometric measurement of glucose 863
K. di Gleria and H. A. O. Hill (Oxford, UK)

The use of rigid organic microemulsions to fix liquid solutions for Mössbauer studies 867
K. Burger, Z. Nemes-Vetéssy (Szeged, Hungary), A. Vértes, Z. Homonnay (Budapest, Hungary), I. Dékány (Szeged, Hungary) and H. Mehner (Berlin-Adlershof, FRG)

The relevance of hydrogen bonding in the mechanism of action of platinum antitumor compounds 873
J. Reedijk (Leiden, Netherlands)

Synthesis of heterocyclic systems by activation of isocyanide, carbonyl, trifluoromethyl and nitrile ligands in platinum(II) and palladium(II) complexes 883
U. Belluco, R. A. Michelin, R. Ros, R. Bertani, G. Facchin, M. Mozzon and L. Zanotto (Padua, Italy)

Author Index 899

Subject Index 901